

**Site of Care**

- ☐ Home Infusion
☐ Infusion Center
☐ Doctor's Office

Fax Referral to (855) 644-3687 or Email: Referrals@sandshealth.com

HYQVIA

(IMMUNE GLOBULIN 10%(HUMAN) WITH RECOMBINANT HUMAN HYLAURONIDASE)

ORDER FORM

PATIENT INFORMATION

Patient Name: _____ DOB: _____
Mobile Number: _____ Patient Weight: _____
Allergies: _____

DIAGNOSIS (Provider must specify)

- ☐ Primary Immunodeficiency (PI), ICD 10: _____
☐ Chronic Inflammatory Demyelinating Polyneuropathy (CIDP), ICD 10: _____
☐ Other: _____

Prior to treatment – ensure the following information is complete and attached with referral:

- ☐ Demographics ☐ Labs and tests supporting diagnosis ☐ Office/progress notes

PRE-MEDICATION

- ☐ Acetaminophen (Tylenol) 500 mg PO ☐ Bendaryl 25 mg PO ☐ Methylprednisolone(Solu-Medrol) _____ mg IVP
☐ Other: _____ Dose: _____ Route: _____
☐ Other: _____ Dose: _____ Route: _____

MEDICATION

MEDICATION	DOSE	ROUTE	FREQUENCY
HyQvia	☐ _____ GM/KG ☐ _____ Grams	☐ Sub Q infusion ☐ Administer as single day infusion ☐ Divide dose over _____ days	every _____ weeks after the initial dose ramp-up*

- ☐ New Start Therapy ☐ Continuation of Therapy Date of last dose (if applicable): _____

LABS/SPECIAL INSTRUCTIONS

Order valid for 1 year from date of signature unless otherwise specified here: _____

PROVIDER INFORMATION

Provider Name: _____ Provider NPI: _____ Contact: _____
Phone: _____ Fax: _____ Email Address: _____
Signature: _____ Date: _____

Ramp up Schedule for Hyqvia

Initial Treatment Interval and Ramp-Up Schedule for PI

For patients previously on another IgG treatment, the first dose should be given approximately 1 week after the last infusion of their previous treatment.

PI: Ramp-up schedule if switching from IVIG

Week	Dose Interval	Dose
1	1st Dose	Total grams x 0.25
2	2nd Dose	Total grams x 0.50
3	NO INFUSION	NO INFUSION
4	3rd Dose	Total grams x 0.75
5	NO INFUSION	NO INFUSION
6	NO INFUSION	NO INFUSION
7	4th Dose	Total grams

PI: Ramp-up schedule if switching from SCIG

Treatment Interval	Dosing Q 4 Weeks	Dosing Q3 Weeks
1st Infusion (Week 1)	Grams x 0.25	Grams x 0.33
2nd Infusion (Week 2)	Grams x 0.5	Grams x 0.67
3rd Infusion (Week 4)	Grams x 0.75	Administer Total Grams
4th Infusion (Week 7)	Administer Total Grams	N/A

Initial Treatment Interval and Ramp-Up Schedule for CIDP

Doses less than or equal to 0.4 g/kg can be administered without ramp-up

Patients must be on stable doses of IVIG for 12 weeks before switching to Hyqvia

Week	Dose Interval	Dose
	Switch from IVIG	
1	No Infusion	
2	1st Dose	Total grams x 0.25
3	2nd Dose	Total grams x 0.25
4	3rd Dose	Total grams x 0.50
6	4th Dose	Total grams x 0.75
9	5th Dose	Total grams

Total grams=total monthly equivalent dose in grams.